

Do Vitamins or Minerals (Apart From Lithium) Have Mood-Stabilizing Effects?

Charles W. Popper, M.D.

Nutritional scientists have been well funded by agribusiness to find ways to deal with factors that interfere with animal health, including aggressive and destructive behavior. When farm animals become “violent”—when pigs start biting each others’ ears and tails, when chickens attack chickens—farmers have learned that the aggressive behavior can be reduced by adding certain minerals and vitamins to their diet, without the need for veterinary intervention.

In 1996, animal nutrition specialist David L. Hardy described this approach to Anthony F. Stephan, whose children had severe treatment-resistant bipolar disorder. Stephan then added similar nutrients to his children’s diet. On the nutritional supplements, both children stabilized clinically and have not needed psychiatric medication for the last 5 years. Hardy and Stephan began advising family members and friends about this nutrient supplement and have now worked with over 2500 psychiatric patients (D. L. Hardy, personal communication, 2001). They also began to collaborate with Bonnie J. Kaplan, Ph.D., a research psychologist at the University of Calgary in Alberta, Canada.

In this issue, Kaplan and colleagues¹ describe an open trial of the first 14 adults with bipolar disorder treated with this nutritional supplement, which consists of a broad range of minerals and vitamins, plus 3 amino acids and several antioxidants. Symptom reductions were clinically noted within 2 weeks and sustained over 6 months of observation. All outcome measures showed significant improvements (55% to 66% symptom reduction), and

a strong effect size ($> .80$) was observed for ratings of depression as well as mania. Most patients could reduce their doses of psychiatric medications, and some patients became stable without any psychiatric medication. Only 2 patients started on new medications that might conceivably have contributed to their stabilization. Even allowing for the usual overestimation of effects in open-label series, these preliminary findings raise interesting questions about nutrition-behavior interactions.

In view of the 50 years of experience with lithium, the notion that minerals can treat bipolar disorder is unsurprising. However, the nutrient supplement studied by Kaplan and colleagues contains no lithium. Might other dietary nutrients have mood-stabilizer properties?

Some may object that a clinical trial of a mixture of ingredients is inherently unscientific: How can one know which ingredient is the active one, whether a smaller number of ingredients will have the same clinical effect, or whether the same ingredients are active in different patients? These questions will become worth pursuing once it has been formally determined whether the mixture, handed down from animal husbandry as a single entity, works in humans. Kaplan’s open-label report justifies her now ongoing controlled study, whose outcome appropriately precedes pursuing questions of mechanism of action and parsimony.

Even so, it is difficult to avoid speculating about possible mechanisms. Might minerals serve as catalysts for enzymes involved in neurotransmitter metabolism, change drug biotransformation, modify membrane receptors or channels, influence second or third messenger systems, or alter gene expression? Lithium, a single cation, has spawned a minor industry of investigations into mechanisms of action,² and the possibilities if there were numerous interacting micronutrients (the collective term for minerals and vitamins) are staggering.

Received Oct. 8, 2001; accepted Oct. 8, 2001. From the Department of Psychiatry, Harvard Medical School, Boston, and McLean Hospital, Belmont, Mass.

Reprint requests to: Charles W. Popper, M.D., McLean Hospital, 115 Mill St., Belmont, MA 02478-1048 (e-mail: Charles_Popper@harvard.edu).

My interest in this nutrient mixture was initially sparked by a case in my clinical practice. A 10-year-old with bipolar disorder was referred for treatment of severe temper tantrums, which had lasted for 2 to 4 hours daily for 4 months. The well-nourished child had no prior psychiatric history or treatment. After 2 days on the Hardy-Stephan nutrient regimen, his tantrums showed significant improvement, with the father-psychiatrist reporting a “complete” absence of outbursts or even irritability at 5 days. After 2 weeks, the available supply of the nutrient supplement was exhausted, and tantrums returned within 48 hours. A similar supplement, containing most of the same ingredients, was then started and produced a moderate improvement, which parents and teachers estimated as 60% of the original effect. When restarted on the original formula, the symptoms were judged to have again responded completely. This naturalistic A-B-A-C-B trial caught my attention because of the full stabilization without psychiatric medications and the absence of observed adverse effects.

I proceeded to cautiously conduct additional trials of the Hardy-Stephan nutrient supplement. Among 22 patients (10 adults, 9 adolescents, 3 preadolescents) who clinically met criteria for bipolar disorder, 19 showed what I judged to be a positive response (2 mild, 7 moderate, 10 marked improvement). Among the 15 patients who were being treated with medications when they began the nutritional supplement, 11 patients have been stable for 6 to 9 months without psychiatric medications. These observations are consistent with Kaplan’s open-label findings, but leave questions of safety unresolved.

Nausea was the main adverse effect in Kaplan’s study. In the larger anecdotal experience of Hardy and Stephan and my limited clinical observations, loose stools and headache were common. Diarrhea, vomiting, flatulence, and agitation were less common. Classical symptoms of mineral or vitamin toxicity were not encountered, but might have emerged with lengthier treatment or more systematic observation.

The Hardy-Stephan supplement contains many nutrients at high doses relative to the Recommended Daily Allowance (RDA), but these RDA levels were primarily established to prevent deficiency disorders in the general population. A broader range of vitamin and mineral functions^{3,4} are considered in formulating the newer daily intake standards, which vary widely because of different criteria for adequacy and different health goals.⁵ Although most ingredients in the nutrient supplement studied by Kaplan and colleagues appear well within safe limits, any multi-ingredient treatment might have toxic effects that cannot be readily predicted from its individual components. Even though it contains only “natural” ingredients and is not under U.S. Food and Drug Administration (FDA) purview, the Hardy-Stephan nutrient supplement should be examined in controlled empirical research—

just as new pharmacologic agents are—to properly assess adverse effects and potential risks.

Psychiatrists do not normally think of vitamins or minerals as modifiers of the effects of psychiatric medications, but the early anecdotal experience with this nutrient supplement suggests that there may be strong *micronutrient-medication interactions*. This mineral-vitamin supplement seems to generally potentiate the clinical properties of psychiatric drugs. Most patients in the Kaplan et al. study could be managed with less medication after the nutrient supplement was added. To avoid medication toxicity, Hardy and Stephan have suggested to patients’ psychiatrists that the doses of psychiatric medications be rapidly reduced shortly after the nutrient supplement is initiated. In my observations, transitioning patients from medications to micronutrients is exceedingly tricky to manage. Introducing micronutrients too quickly can increase the adverse effects of medications, including agitation, while withdrawing psychiatric medications too quickly can result in symptom exacerbation. Often, both increased adverse effects and symptom resurgence are seen at once. Much more data are needed about how to “transition” patients who are currently taking psychiatric medications. Although it appears less difficult to treat medication-naïve patients (such as my 10-year-old patient), the transition from psychiatric medications to micronutrients can require genuine technical savvy—even for patients who have not taken such medications for several weeks or months. Clinicians who mistakenly approach these new findings as encouragement to combine micronutrients with psychiatric medications may find that they have stepped into a serious quagmire.

Although health food advocates have made numerous claims without scientific documentation, nutritional influences on mental illness have received considerable research attention,⁶ some of which is quite rigorous and promising. Andrew Stoll’s research on omega-3 fatty acids for bipolar disorder^{7,8} and Eugene Arnold’s work on omega-6 fatty acids for attention deficit disorder⁹⁻¹¹ suggest that these nutrients might themselves be therapeutic. Other micronutrients (calcium, chromium) and macronutrients (inositol, amino acids) have also shown some potential for influencing mood disorders.¹²⁻¹⁵ Arnold’s recent findings suggest that relative zinc deficiency might explain why some patients with ADHD do not show a more robust response to psychostimulants.⁹

Several large-scale double-blind placebo-controlled studies of RDA or high-dose multivitamin regimens in adults have reported improved scores of mood and cognition^{16,17} as well as anxiety and somatic symptoms.¹⁸ A recent review reported that 10 of 12 randomized double-blind placebo-controlled studies found that multi-micronutrient regimens can improve cognitive functioning in children.¹⁹ However, high-dose multivitamin treatments have not been found to be effective in children with ADHD or learning disorders.^{11,20}

Speculatively, there is a great deal to learn about minerals and vitamins in psychiatric pathophysiology and psychopharmacology. It is intriguing that many of the enzymes proposed as target sites of lithium action are metalloenzymes that are noncompetitively inhibited by lithium, which probably acts by displacing the divalent cation.²¹ Why would supplementation with divalent cations have a therapeutic effect? Do we know enough about complex intracellular regulatory interactions to answer this question? Individual micronutrients will need to be examined in combination with lithium, but it is unlikely that the physiology will be so simplistic and dyadic. The groundbreaking approach of examining several nutrient ingredients at once, while a violation of our usual tenets of investigation, may present an opportunity to examine how micronutrients might operate in concert.

Assessing the safety and efficacy of multinutrient formulas will require considerable research. Developing dose-effect curves for each micronutrient, and examining all possible combinations of micronutrients, will be a horrifically large task.²² Many years will be required to arrive at a formula that is optimal for the general population, but it is more likely there will eventually be different formulas whose safety and efficacy are optimized to the metabolic requirements of treating different disorders, different individuals, different ages, and different comorbid health situations.

The possibility of a nutritional alternative to drug treatment may raise hope and carry the risk of igniting public interest beyond reasonable bounds. Some patients may find it difficult to wait for nutrient supplements to be examined in humans for both efficacy and safety in controlled trials, and clinicians will need to help patients keep their enthusiasm from pushing usage beyond its scientific basis. Similarly, clinicians will want to think twice if they are tempted to forge ahead with empirical trials of this novel treatment with possible unknown risks and unproven benefit. Some physicians, presuming a low risk of toxicity, may reason that it is judicious to allow some drug-naïve patients to proceed with empirical trials of nutrient supplements for a few weeks before committing them to treatment with psychiatric medications. Most clinicians will want the reassurance of systematic safety data before beginning to examine its effects in patients. Clinicians will hopefully minimize risks of nutrient-medication interactions by initially restricting their trials to patients who have not recently used psychiatric medications.

If Kaplan and colleagues' preliminary findings are confirmed in controlled research, and if safety studies are favorable, what then? What if some psychiatric patients could be treated with inexpensive vitamins and minerals rather than expensive patented pharmaceuticals? or what if some doses of psychiatric drugs could be reduced by the

concurrent use of nutrients? The economic implications, for individual patients and for the pharmaceutical industry, are difficult to overlook. For now, micronutrient treatments and other nutritional approaches remain in a very early stage of scientific investigation. Depending on how this line of research develops, clinicians and researchers may need to rethink the traditional bias against nutritional supplementation as a potential treatment for major psychiatric disorders.

REFERENCES

1. Kaplan BJ, Simpson JSA, Ferre RC, et al. Effective mood stabilization with a chelated mineral supplement: an open-label trial in bipolar disorder. *J Clin Psychiatry* 2001;62:936-944
2. Shalubina A, Agam G, Belmaker RH. The mechanism of lithium action: state of the art, ten years later. *Prog Neuropsychopharmacol Biol Psychiatry* 2001;25:855-866
3. Mertz W. A perspective on mineral standards. *J Nutr* 1998;128(suppl 2): 375S-378S
4. Cashman KD, Flynn A. Optimal nutrition: calcium, magnesium and phosphorus. *Proc Nutr Soc* 1999;58:477-487
5. Mertz W. Three decades of dietary recommendations. *Nutr Rev* 2000;58: 324-331
6. Werbach MR. *Nutritional Influences on Mental Illness: A Sourcebook of Clinical Research*. 2nd ed. Tarzana, Calif: Third Line Press; 1999
7. Stoll AL, Locke CA, Marangell LB, et al. Omega-3 fatty acids and bipolar disorder: a review. *Prostaglandins Leukot Essent Fatty Acids* 1999;60: 329-337
8. Stoll AL, Severus E, Freeman MP, et al. Omega 3 fatty acids in bipolar disorder: a preliminary double-blind, placebo-controlled trial. *Arch Gen Psychiatry* 1999;56:407-412
9. Arnold LE, Pinkham SM, Votolato N. Does zinc moderate essential fatty acid and amphetamine treatment of attention-deficit/hyperactivity disorder? *J Child Adolesc Psychopharmacol* 2000;10:111-117
10. Arnold LE. Alternative treatments for adults with attention-deficit hyperactivity disorder (ADHD). In: Wasserstein J, Wolfe LE, Lefever FF, eds. *Adult Attention Deficit Disorders: Brain Mechanisms and Life Outcomes*. New York, NY: New York Academy of Sciences; 2001:310-341
11. Arnold LE. Treatment alternatives for attention-deficit/hyperactivity disorder. In: Jensen PS, Cooper J, eds. *Diagnosis and Treatment of ADHD: An Evidence-Based Approach*. Washington, DC: American Psychiatric Press. In press
12. Levy NA, Janicak PG. Calcium channel antagonists for the treatment of bipolar disorder. *Bipolar Disord* 2000;2:108-119
13. McLeod MN, Golden RN. Chromium treatment of depression. *Int J Neuropsychopharmacol* 2000;3:311-314
14. Chengappa KN, Levine J, Gershon S, et al. Inositol as an add-on treatment for bipolar depression. *Bipolar Disord* 2000;2:47-55
15. Van der Does AJ. The effects of tryptophan depletion on mood and psychiatric symptoms. *J Affect Disord* 2001;64:107-119
16. Benton D, Fordy J, Haller J. The impact of long-term vitamin supplementation on cognitive functioning. *Psychopharmacology (Berl)* 1995;117: 298-305
17. Benton D, Haller J, Fordy J. Vitamin supplementation for 1 year improves mood. *Neuropsychobiology* 1995;32:98-105
18. Carroll D, Ring C, Suter M, et al. The effects of an oral multivitamin combination with calcium, magnesium, and zinc on psychological well-being in healthy young male volunteers: a double-blind placebo-controlled trial. *Psychopharmacology (Berl)* 2000;150:220-225
19. Benton D. Micro-nutrient supplementation and the intelligence of children. *Neurosci Biobehav Rev* 2001;125:297-309
20. Arnold LE. Treatment alternatives for attention-deficit hyperactivity disorder (ADHD). *J Attn Disord* 1999;3:30-48
21. Phiel CJ, Klein PS. Molecular targets of lithium action. *Annu Rev Pharmacol Toxicol* 2001;41:789-813
22. Pryor WA. Vitamin E and heart disease: basic science to clinical intervention trials. *Free Radic Biol Med* 2000;28:141-164

Effective Mood Stabilization With a Chelated Mineral Supplement: An Open-Label Trial in Bipolar Disorder

Bonnie J. Kaplan, Ph.D.; J. Steven A. Simpson, Ph.D., M.D.;
Richard C. Ferre, M.D.; Chris P. Gorman, M.D.;
David M. McMullen, M.D.; and Susan G. Crawford, M.Sc.

Background: To determine in open trials the therapeutic benefit of a nutritional supplement for bipolar disorder.

Method: The sample consisted of 11 patients with DSM-IV–diagnosed bipolar disorder aged 19 to 46 years, who were taking a mean of 2.7 psychotropic medications each at study entry. Three additional patients dropped out prematurely. The intervention is a broad-based nutritional supplement of dietary nutrients, primarily chelated trace minerals and vitamins, administered in high doses. At study entry and periodically thereafter, patients were assessed with the Hamilton Rating Scale for Depression (HAM-D), the Brief Psychiatric Rating Scale (BPRS), and the Young Mania Rating Scale (YMRS).

Results: For those who completed the minimum 6-month open trial, symptom reduction ranged from 55% to 66% on the outcome measures; need for psychotropic medications decreased by more than 50%. Paired *t* tests revealed treatment benefit on all measures for patients completing the trial: HAM-D mean score at entry = 19.0, mean score at last visit = 5.4, *t* = 5.59, *df* = 9, *p* < .01; BPRS mean score at entry = 35.3, mean score at last visit = 7.4, *t* = 2.57, *df* = 9, *p* < .05; YMRS mean score at entry = 15.1, mean score at last visit = 6.0, *t* = 4.11, *df* = 9, *p* < .01. The effect size for the intervention was large (> .80) for each measure. The number of psychotropic medications decreased significantly to a mean ± SD of 1.0 ± 1.1 (*t* = 3.54, *df* = 10, *p* < .01). In some cases, the supplement replaced psychotropic medications and the patients remained well. The only reported side effect (i.e., nausea) was infrequent, minor, and transitory.

Conclusion: Some cases of bipolar illness may be ameliorated by nutritional supplementation. A randomized, placebo-controlled trial in adults with bipolar I disorder is currently underway, as well as open trials in children.

(*J Clin Psychiatry* 2001;62:936–944)

Received Feb. 27, 2001; accepted June 11, 2001. From the Faculty of Medicine, University of Calgary (Drs. Kaplan and Simpson) and the Alberta Children's Hospital (Dr. Kaplan and Ms. Crawford), Calgary, Alberta, Canada. Drs. Gorman and McMullen are in private practice in Calgary, Alberta, and Dr. Ferre is in private practice in Salt Lake City, Utah.

Supported in part by the Alberta Children's Hospital Foundation and the Alberta Science and Research Authority, Calgary, Alberta; and Evince International, Farmington, Utah (who provided the E.M.Power+ supplement free of charge).

Presented in part at the 50th Annual Meeting of the Canadian Psychiatric Association in October 2000 and at the meeting of the Society of Biological Psychiatry in May 2001.

We thank Dr. Catherine Field, Mr. Anthony Stephan, Mr. David Hardy, and Dr. Charles Popper for their support and consultation.

Reprint requests to: Bonnie J. Kaplan, Ph.D., Alberta Children's Hospital, 1820 Richmond Rd. S.W., Calgary, AB T2T 5C7, Canada (e-mail: kaplan@ucalgary.ca).

Solid scientific research shows that many dietary nutrients, including minerals and vitamins, are essential for normal brain function. For instance, deficient levels of various B vitamins are related to pathologic brain and behavior disorders ranging from Korsakoff's syndrome to pellagra. Recent work on folic acid (vitamin B₉) suggests that low levels may be associated with depressive symptomatology and poor response to antidepressant medication.^{1,2}

Less is known about the role of trace elements, but there is considerable evidence that these too may be essential for normal brain function. Zinc provides a good example. Most of the excitatory neurons of the cerebral cortex have glutamate as their primary transmitter. One type of glutaminergic neuron accumulates zinc within vesicles at axon terminals and releases it into the synapse upon firing.³ The precise roles of zinc in synaptic function are not known, although its presence is certain,⁴ and there are zinc-binding sites on one subset of glutamate receptor called the NMDA (*N*-methyl-D-aspartate) receptor.⁵ Zinc, copper, and magnesium all appear to play important modulatory roles in controlling the NMDA receptor,^{6–8} which has been implicated in various forms of cortical plasticity, including learning.⁹ It is possible, then, that decreased levels of some minerals in the brain may produce abnormal NMDA-mediated plasticity and subsequent abnormalities in behav-

See also Commentary on page 933.

ior. Other trace elements such as magnesium and copper^{6,10} are also thought to have receptor sites in the cortex and might also presumably affect normal cortical functioning.

Recent studies have shown low plasma levels of zinc and other minerals in people with mood and behavior disorders. Maes and colleagues^{11,12} found lower serum zinc levels in 48 unipolar depressed patients in comparison to 32 volunteers with normal mental health. Walsh and colleagues¹³ evaluated the copper and zinc levels in 135 assaultive, incarcerated males in comparison to controls. They found an inverse relationship between zinc and the seriousness of the behavior, ranging from verbal assault to aggravated and violent assault: the lower the zinc, the greater the rate of violent behavior.

Toren and colleagues¹⁴ reported lower serum zinc levels in 43 children with attention-deficit/hyperactivity disorder (ADHD) than in comparable controls. A study of 18 boys with ADHD who underwent a placebo-controlled trial of amphetamine treatment found that higher baseline hair zinc levels predicted a better behavioral response to the stimulant medication.¹⁵ Subsequent analyses of the same data, but employing a more thorough determination of zinc levels from 3 measures (hair, red cell, and urine), confirmed the initial finding: those children with adequate zinc were better stimulant responders than those with borderline zinc levels or those who were zinc-deficient.¹⁶

Studies of other minerals have yielded some interesting findings. Calcium imbalance caused by hyperparathyroidism has long been known to affect anxiety, depression, and cognitive function,^{17,18} and in fact, calcium balance in general has been shown to play an important role in mood disorders.¹⁹ A recent study reported significantly decreased bone mineral density in 24 women with current major depressive disorders or with a past history of depression,²⁰ and women with osteoporosis exhibited elevated depression scores on a symptom inventory.²¹ Significant iron deficiency results in mental and behavioral symptoms such as irritability, aggression, or mental retardation.^{22,23} Because iron-deficient rats have low dopamine D₂ receptor levels,²⁴ Weiser and colleagues²⁵ looked at 26 medication-free schizophrenics in comparison to controls and found significantly low serum iron. Others have reported that both the cerebrospinal fluid and serum of people with schizophrenia were low in magnesium.²⁶

The evidence supporting the importance of minerals and vitamins in central nervous system functioning provided some scientific support for our interest in studying a supplement that is gaining considerable clinical attention in Canada and in many areas of the United States. This supplement was developed over several years by some families in Alberta and eventually was manufactured and distributed from Utah.* It consists of 36 ingredients, primarily chelated minerals. The formulation was based in part on agricultural knowledge of the treatment of stress and mood problems in cattle and hogs: transport and han-

dling of livestock have been shown to have a particularly large impact on chloride, potassium, calcium, and magnesium, thus altering normal electrolyte balance and overall health.²⁷ Described herein are the data collected in open trials of the first, consecutive, and unselected 11 cases of bipolar disorder in adults evaluated while taking this supplement and followed for at least 6 months, testing the hypothesis that a broad-based nutritional supplement that emphasizes trace minerals would help stabilize mood.

METHOD

The Conjoint Health Research Ethics Board of the University of Calgary, Calgary, Alberta, Canada, approved the protocol for an open case series in adults with bipolar disorder, using the dietary supplement E.M.Power+. The experimental nature of the work was carefully described to each patient, who then signed a consent form.

The overall sample for this open-label case series consisted of 14 adults (10 men and 4 women). The 3 patients who dropped out were female, aged 24, 35, and 45 years. The dropout patients took the supplement for at least 3 weeks (range, 3–10 weeks). The sample of completers consisted of 10 men and 1 woman, aged 19 to 46 years (mean \pm SD = 29.4 \pm 10.7 years). Six cases fulfilled DSM-IV criteria for bipolar I, 4 fulfilled criteria for bipolar II, and 1 fulfilled criteria for bipolar disorder not otherwise specified (NOS). Patients' psychiatrists confirmed their clinical diagnoses using the Structured Clinical Interview for DSM-IV Axis I Disorders (the SCID-I).²⁸ The time since diagnosis for each patient is listed in Table 1, along with their medication histories.

Patients were excluded if they were pregnant, trying to get pregnant, or lactating; if they had untreated or unstable thyroid disease; if they had received ECT within the previous 6 months; if their psychotropic medications had been altered within the previous 4 weeks; if they were currently taking antibiotics (which may disrupt intestinal absorption of nutrients); if they had any abnormality of metabolism (renal disease, hemochromatosis, hyperlipidemia, etc.); if there was evidence of substance abuse in the previous 6 months; if they suffered from dementia, mental retardation, or any neurologic disorder affecting language or comprehension; if they suffered from any other unstable medical condition; or if they had begun any other new treatment recently such as cognitive psychotherapy. Patients were not excluded for comorbid diagnoses: 1 patient had obsessive-compulsive disorder, 1 had dysthymic disorder, and 1 had ADHD. Participants were

*After several years of development by the Synergy Group of Canada/Truehope Institute, in April 2000, the formulation of the supplement was finalized and named E.M.Power+ ("essential mineral" power). A list of ingredients for 1/4 the full adult dose can be found in the Appendix.

Table 1. Characteristics of 14 Patients With Bipolar Disorder

| Patient | Age (y), Gender, Diagnosis | Years Since Diagnosis | Past Medications | ECT?/Hospitalizations? |
|---------|----------------------------------|-----------------------------|---|------------------------|
| 1 | 21, M, bipolar I | 2 | Benzotropine, bupropion, buspirone, clonazepam, chlorpromazine, dextroamphetamine, diazepam, fluoxetine, imipramine, lithium, methylphenidate, nortriptyline, paroxetine, risperidone, sertraline, trazodone, valproic acid, zolpidem | No/No |
| 2 | 19, M, bipolar I | 3 | Carbamazepine, clomipramine, clonazepam, desipramine, fluoxetine, fluvoxamine, haloperidol, imipramine, lamotrigine, lithium, nortriptyline, paroxetine, quetiapine, risperidone, sertraline, thioridazine, valproic acid | Yes/Yes |
| 3 | 23, M, bipolar NOS | 4 | Bupropion, dextroamphetamine, fluoxetine, pemoline, sertraline, venlafaxine | No/No |
| 4 | 20, M, bipolar I | 3 | Desipramine, fluoxetine, imipramine, lithium, methylphenidate, paroxetine, sertraline, valproic acid, venlafaxine | No/No |
| 5 | 21, M, bipolar II | 3 | Clonazepam, fluoxetine, paroxetine, zolpidem | No/Yes |
| 6 | 45, M, bipolar II | 3 | Bupropion, dextroamphetamine, fluoxetine, lithium, methylphenidate, thioridazine, trifluoperazine, valproic acid, venlafaxine | No/Yes |
| 7 | 34, M, bipolar I | 16 | Alprazolam, clonazepam, desipramine, fluoxetine, lithium, risperidone, sertraline, trazodone, valproic acid, venlafaxine, zolpidem | No/No |
| 8 | 46, M, bipolar I | 7 | Bupropion, clonazepam, clozapine, fluoxetine, gabapentin, lamotrigine, lithium, olanzapine, quetiapine, valproic acid | Yes/Yes |
| 9 | 21, M, bipolar I | 3 | Paroxetine, valproic acid | No/Yes |
| 10 | 31, M, bipolar II | 2 | Chlorpromazine, citalopram, dextroamphetamine, fluvoxamine, gabapentin, lamotrigine, levothyroxine, lithium, lorazepam, methotrimeprazine, nefazodone, olanzapine, sertraline, valproic acid, venlafaxine | No/Yes |
| 11 | 42, F, bipolar II | 19 | Carbamazepine, lithium, sertraline, trifluoperazine | Yes/Yes |
| 12 | 35, F, bipolar I | 18 | Clonazepam, lithium carbonate, lorazepam, olanzapine, paroxetine, risperidone, topiramate, valproic acid, zopiclone | No/Yes |
| 13 | 45, F, bipolar I | 20 | Amitriptyline, citalopram, doxepin, fluoxetine, imipramine, lithium carbonate, lorazepam, methotrimeprazine, nefazodone, trazodone, trimipramine maleate, venlafaxine | Yes/Yes |
| 14 | 24, F, bipolar II | 5 | Bupropion, carbamazepine, chlorpromazine, clonazepam, fluoxetine, lithium carbonate, nefazodone, paroxetine, risperidone, sertraline, valproic acid | No/Yes |

asked to stop all other nutritional supplementation at study entry.

The intervention consisted of the mineral and vitamin formulation in capsule form. The full dose requires the ingestion of 32 capsules per day because of the bulk of chelated minerals. Patients were started immediately on the full dose on the first treatment day and were allowed to distribute the capsules into 4 doses across the day.

Patients were assessed by their own psychiatrists at study entry and during the intervention with the following measures: the Hamilton Rating Scale for Depression (HAM-D),²⁹ the Young Mania Rating Scale (YMRS),³⁰ and the Brief Psychiatric Rating Scale (BPRS).³¹ These 3 outcome measures were employed at study entry and at each

visit, along with an Adverse Events Form used to monitor side effects. Patients were seen by their psychiatrists once a week for the first 4 weeks of the open trial. Appointments were decreased to a monthly schedule if a patient's score on the HAM-D decreased to less than 8 (a HAM-D score of about 14–17 is generally considered to indicate clinically important levels of depression)³² or if the score on the BPRS decreased to less than 6 (a BPRS score of 14–20 is generally thought to indicate clinically meaningful levels of psychiatric symptomatology).^{33,34} Patients attended at least 7 appointments with their psychiatrists during the intervention (range, 7–18 appointments).

Psychiatrists were permitted to manage their patients as usual, using their clinical judgment regarding care. This

Table 2. Medications Before and During Treatment With a Chelated Mineral Supplement

| Patient | Medications at Study Entry | Time on Supplement (wk) | Current Medications |
|---------|---|-------------------------|--|
| 1 | Clonazepam, lithium, nortriptyline, risperidone, zolpidem | 40 | Clonidine, lamotrigine, olanzapine, quetiapine |
| 2 | Carbamazepine, clonazepam, lamotrigine, propranolol, quetiapine, sertraline | 59 | Lamotrigine |
| 3 | None | 41 | None |
| 4 | None | 58 | None |
| 5 | Paroxetine | 28 | None |
| 6 | Fluoxetine, lithium, methylphenidate, venlafaxine | 86 | Nortriptyline |
| 7 | Clonazepam, zolpidem | 26 | Zolpidem |
| 8 | Clonazepam, lamotrigine, lithium, quetiapine | 34 | Quetiapine |
| 9 | Paroxetine, valproic acid | 45 | None |
| 10 | Lithium, lorazepam, sertraline, valproic acid | 40 | Lithium |
| 11 | Carbamazepine, sertraline | 26 | Sertraline |
| 12 | Risperidone, topiramate | 6 | None |
| 13 | Clonazepam, moclobemide, valproic acid | 10 | Clonazepam, moclobemide |
| 14 | Citalopram, lorazepam, topiramate | 3 | Citalopram, lorazepam, quetiapine, topiramate |

included adjustment of concurrent psychiatric medications as they saw fit. They were, however, also given instructions based on the developers'/distributor's suggestions on managing expected supplement-medication interactions. The instruction was that medication reductions should be considered in the event of interactions.

Fourteen patients began the protocol and 11 completed a minimum of 6 months; as shown in Table 2, we were able to follow most for much longer (range, 6–21 months). The reasons that patients dropped out of the study (N = 3) were varied. In 1 case that was monitored by someone outside the research team, the psychiatrist violated the protocol and did not monitor the supplement-medication interactions as required. This patient took the supplement for 3 weeks and became very agitated and tearful. The psychiatrist admitted the patient to hospital and discontinued the supplement rather than altering medication. In the other 2 cases, definite symptom improvement was demonstrable on the daily and periodic data collection systems. Both patients became noncompliant, however, and were removed from the protocol because of refusing to follow their research psychiatrists' recommendations. In 1 of the cases, the patient took the supplement for 6 weeks as part of this study and was able to eliminate both of the medications that she was taking initially. This patient is continuing to take the supplement on her own (apparently with considerable success). She did, however, refuse to submit to a final assessment for this study, because of conflicts with psychiatrists on the research team.

Statistical Analysis

The outcome variables were analyzed using paired t tests to assess the difference between scores at study entry com-

pared with scores at the last study visit. Two sets of analyses were conducted: intent-to-treat analysis (N = 14) and analysis of patients who completed the open trial (N = 11). The effect size for each outcome variable was calculated by dividing the absolute value of the mean of the paired difference by the standard deviation of the difference.

RESULTS

Ten patients had been hospitalized at least once, of whom 4 had prior electroconvulsive therapy (see Table 1). The mean number of psychotropic medications tried prior to study entry was 10 (range, 2–18), and mean years of prior treatment were 6.7 (range, 2–20).

Table 2 lists the psychotropic medications at entry into the trial, the number of weeks each patient has taken the supplements thus far, and the medications at the most recent visit. In some cases, the lack of medication at study entry is misleading. For instance, patient 4 had taken many medications for the 3 years prior to this study (as shown in Table 1). None had been effective, so a month prior to entering this study he had chosen to discontinue all psychotropic medications. At entry, the mean \pm SD number of medications prescribed for this sample was 2.7 ± 1.8 .

Symptom Reduction

Intent-to-treat analysis. A series of paired t tests with all 14 subjects revealed statistically significant decreases in scores over the course of the study on the HAM-D and YMRS (Table 3). There was also a trend for lower scores on the BPRS.

Analysis of completers. The same analyses were repeated on the 11 patients who completed the open trial.

Table 3. Outcome Measures at Study Entry and at Last Visit^a

| Outcome | Entry Visit | | Last Visit | | Full Sample | | | Restricted Sample ^b | | |
|--------------------------|-------------|------|------------|------|-------------|----|-------|--------------------------------|----|-------|
| | Mean | SD | Mean | SD | Paired t | df | p | Paired t | df | p |
| Intent-to-treat (N = 14) | | | | | | | | | | |
| HAM-D | 17.8 | 8.1 | 9.2 | 9.0 | 2.64 | 12 | < .05 | | | |
| BPRS | 30.8 | 30.1 | 10.2 | 11.0 | 2.02 | 11 | < .10 | | | |
| YMRS | 15.9 | 11.9 | 6.0 | 6.4 | 3.73 | 11 | < .01 | | | |
| Completers only (N = 11) | | | | | | | | | | |
| HAM-D | 19.0 | 8.0 | 5.4 | 5.6 | 5.59 | 9 | < .01 | 4.99 | 8 | < .01 |
| BPRS | 35.3 | 31.2 | 7.4 | 8.6 | 2.57 | 9 | < .05 | 2.34 | 8 | < .05 |
| YMRS | 15.1 | 9.8 | 6.0 | 6.9 | 4.11 | 9 | < .01 | 3.61 | 8 | < .01 |

^aAbbreviations: BPRS = Brief Psychiatric Rating Scale, HAM-D = Hamilton Rating Scale for Depression, YMRS = Young Mania Rating Scale.

^bRestricted sample consists of 10 patients who completed the study, omitting patient 1.

Table 4. Change in Outcome Measures from Study Entry to Last Visit by Subject

| Patient | HAM-D | | BPRS | | YMRS | |
|-----------------|-------|------|-------|------|-------|------|
| | Entry | Last | Entry | Last | Entry | Last |
| 1 ^a | 28 | 14 | 40 | 15 | 27 | 16 |
| 2 ^a | 25 | 2 | 54 | 2 | 12 | 1 |
| 3 ^a | 18 | 2 | 38 | 2 | 22 | 3 |
| 4 ^a | 21 | 1 | 113 | 3 | 6 | 3 |
| 5 | 16 | ... | 23 | 21 | 18 | 16 |
| 6 ^a | 17 | 3 | 29 | 2 | 15 | 0 |
| 7 | 11 | 6 | 10 | 22 | 25 | 15 |
| 8 ^a | 25 | 16 | 20 | 7 | 24 | 6 |
| 9 | 21 | 8 | ... | ... | 0 | 0 |
| 10 ^a | 23 | 0 | 25 | 0 | 2 | 0 |
| 11 | 1 | 2 | 1 | 0 | 0 | 0 |
| 12 | 9 | 15 | 12 | 34 | 38 | 9 |
| 13 | 24 | 28 | 20 | ... | 2 | ... |
| 14 | 8 | 22 | 5 | 14 | 2 | 3 |

^aPatient is a responder (as indicated by $\geq 20\%$ reduction on all 3 measures).

Paired t tests revealed significant decrements in scores from study entry to the most recent visit on all 3 outcome measures assessing depression (HAM-D), mania (YMRS), and general psychiatric status (BPRS) (see Table 3). The effect size was large for each of the 3 outcome measures ($> .80$). These patients have now been followed for a mean \pm SD of 43.9 ± 18.0 weeks (range, 26–86 weeks).

The mean symptom reduction was 55% on the HAM-D, 60% on the BPRS, and 66% on the YMRS (Table 4). To evaluate the number of patients showing a clinically significant response to the supplement, we defined as a responder any patient who exhibited at least a 20% decrease on all 3 outcome measures. Two patients (patient 5 and patient 9) were excluded because of missing data, and 1 patient (patient 11) because she was asymptomatic at study entry, so that a 20% decrease was not possible. Of the remaining 8 patients, 7 met criteria for responder status (Table 4). The remaining patient (patient 7) met responder criteria for 2 of the 3 measures. Although his BPRS score actually increased considerably from 10 to 22, an examination of his interview data revealed that this was not due to the presence of any psychotic or manic symptoms. His symptoms were primarily

guilt, anxiety, and tension, which his psychiatrist reports were due to a profound reaction to the discovery that his family had found out that he was significantly in debt from a gambling addiction.

Medication Reduction in Completers

The number of psychotropic medications was reduced by more than 50% in these patients after they began taking the supplement, from a mean of 2.7 ± 2.0 per patient at study entry to 1.0 ± 1.1 at the most recent visit ($t = 3.54$, $df = 10$, $p < .01$). The protocol allowed psychiatrists to change medications freely as part of their patient management. Two patients were medication-free at the start of the open trial and at their last visit: patient 3 was medication-free during the entire study period; patient 4 tried 2 medications during the open trial but was medication-free at the start of the study and again by the last study visit. All other patients were on different medication regimens at the start and finish. The medication regimen of 1 patient (patient 1) was especially different at the end of the trial compared with study entry, so that the patient's results were discarded and the data were reanalyzed. As shown in Table 3, the results from this restricted sample were similar to the overall results.

In 2 cases, the supplement has replaced psychoactive medications and the patients have remained well. In general, patients have reported feeling the first effect of the supplement within 2 weeks. Medications were decreased gradually as the supplement began to influence the patient's mood.

Side Effects

The only side effect reported was nausea, especially if patients forgot to take the supplement with food. In 2 cases the nausea was sustained over several days. Patient 9 chose to increase his food intake with his pills and briefly use antinausea medication to cope with the problem; the nausea of patient 10 was alleviated by dose reduction and a gradual increase up to full dose.

Despite the challenge of taking 32 pills daily, the patients reported that the supplement was easy to tolerate. Also, some patients reported being able to distinguish

their subjective reactions to the supplement (which they described as feeling more normal) from the benefit they have derived from psychotropic medication treatment (described as symptom reduction or masking).

DISCUSSION

We have been exploring the possible clinical benefit of a broad-based nutritional supplement consisting of 36 items, primarily chelated minerals. The data reported here provide the first, preliminary scientific validation of the supplement's efficacy and suggest that further research is warranted.

Although much additional research is needed (to replicate these findings, to address the mechanism of action, etc.), our preliminary data and the general clinical experience of psychiatrists who are monitoring patients in our trials indicate that the supplement has a beneficial psychotropic effect and is not acting in only an adjunctive manner. Consequently, the largest challenge in dealing with the supplement is the interaction with psychiatric medications. The observation made by many patients and clinicians who have used this preparation is that the supplement interacts with psychiatric medications. The distributor recommends decreasing psychiatric medications in this situation, and despite significant concerns about safety, we have found that this seems to be a reasonable approach.

In this case series, the diagnoses of bipolar disorder in these 11 patients were made clinically by psychiatrists and confirmed by the SCID diagnostic interview. Patients were treated with the nutritional supplement on an open-label protocol and monitored systematically for a minimum of 6 months (range, 6–21 months). The changes in all 3 outcome measures were statistically and clinically significant. As a group, it was possible for the patients to be clinically managed on fewer psychotropic medications (more than a 50% reduction). Individually, also, all 11 patients benefited from the supplement. In the 1 patient who was asymptomatic at study entry (patient 11), whose benefit cannot be demonstrated as a symptom reduction, a decrease in number of medications was achieved without a significant elevation in symptoms.

There are weaknesses in this case series. First, purely by chance, 10 of these patients were men, thus hindering generalizations to women. Second, as in any case series, there is a weakness in not having a placebo control. We would point out that these patients had been exposed previously to many interventions, and their positive expectancy of benefit did not work with these prior efforts. Also, it would be surprising for a placebo effect to be sustained as long as 6 months. Nevertheless, a randomized, placebo-controlled trial has been funded and is currently underway. A third potential source of bias is from the psychiatrists themselves. As in any open-label study, unblinded assessments can result in exaggerated results. In this particular

case, though, it would be fair to say that the majority of the psychiatrists expected to see no benefit from nutritional supplementation, so perhaps this form of bias was not a major influence on our data. The use of concurrent psychiatric medication was a fourth weakness in this case series: the changes in medications made by the psychiatrists as part of normal clinical care make it difficult to attribute symptom changes specifically to the nutritional supplement. Our current randomized trials are evaluating patients who are unmedicated.

It would be a mistake to assume that the breadth of the intervention was a weakness. At present, there is not enough knowledge available to hypothesize that one or several of the ingredients has any special effectiveness in treating bipolar disorder. The selection of a broad-based mineral and vitamin supplement was based on an approach of inclusiveness rather than premature focus or closure, as well as a strategy of attempting to maximize the chances that some clinical effect might be found. There have been many failed attempts to treat mental illness with a single ingredient or group of nutrients, and also possibly some limited successes.^{35,36} Given the complexity of the mammalian central nervous system and homeostatic mechanisms, it is reasonable that a very broad approach to nutritional supplementation might actually be appropriate.

This initial broad-based strategy allows a series of future studies that might focus on particular groups of nutrients or eventually a selective series of active components. However, at this time it is not known whether a subset of components will be effective or whether a mixture of a broad range of nutrients will be required to maximize clinical effectiveness. To those who are concerned that it is not known which of the 36 nutrients is "the important one," we would say that the likelihood of finding a single effective ingredient is very small. Our future research will, in fact, be attempting to decrease the number of ingredients, in part to reduce the bulk of the supplement. For now, the complexity of the formulation should not detract from the treatment findings and, indeed, might be one of its strengths. Similarly, the doses of the individual ingredients of the nutritional supplement were quite high, again to maximize the chances of observing a clinical response, with the expectation that it may be possible in the future to lower the doses as a result of subsequent studies.

These data also have many strengths. First, the 11 cases in the completer analysis were unselected: they were the first 11 patients with bipolar disorder who were assessed systematically and followed for a minimum of 6 months. Also, in spite of receiving optimal psychiatric care, these were patients who were still symptomatic (or, in the case of patient 11, uncomfortable with the side effects of their medications). No problematic adverse effects were encountered from the nutrient supplement. Patients generally reported a subjective sense of improved well-being when taking the supplement, and several patients described this

well-being as feeling more normal than what they had experienced with psychotropic medication.

Safety

Vitamin toxicity is a serious consideration when high doses of these substances are administered on a chronic basis. Patients with any known metabolic disorder (Wilson's disease, hemochromatosis, etc.) were excluded from this protocol. However, the assurance of safety of this supplement for any healthy patient is not something that can be precisely stated at this time. Prospective studies to establish an intake associated with toxicity of vitamins and minerals are not studies that have been, or even ethically can be, conducted. Data can be collected only from case studies of individuals who medicate themselves, or have ingested large quantities, and toxicity predicted to a very limited extent from animal studies. Additional problems in trying to set a level of intake arise from the knowledge that the bioavailability of some vitamins and most minerals depends on the rest of the foods in the diet, the presence of other dietary nutrients, the nutritional status of the individuals (absorption is significantly reduced when status is inadequate), and an individual's genetic makeup.

The levels of several ingredients might raise some questions regarding potential toxicity. Vitamin A is present in the supplement at a daily intake of 13,332 IU, or about 4 times the recommended daily intake, but chronic toxic reactions would not be expected at dosages less than 100,000 IU/day in adults with normal liver function.³⁷ Pregnant and lactating women are excluded from our protocols because of uncertainty about fetal exposure. Vitamin D in the supplement is present at a daily level of 1600 IU (40 µg) in the form of cholecalciferol, and in adults vitamin D toxicity usually does not occur unless intakes exceed 50,000 IU/day.³⁸ Vitamin B₁₂ is provided in the supplement at 1000 µg/day, but there has been no reported toxicity associated with intakes of vitamin B₁₂ at levels 1000 times the recommended daily intake of 6 µg/day; this is probably because the absorption decreases as body stores of vitamin B₁₂ increase. Animal studies have demonstrated no adverse effects of vitamin B₁₂, even when administered by injection in very large doses.³⁹ Selenium in the supplement provides 400 µg/day in the form of a selenium amino acid chelate; studies have indicated that there are no clinical signs of selenosis at intakes of 853 µg/day.⁴⁰ Chromium is supplied in the supplement at 1000 µg/day. As there are no reported case studies available on toxicity, its ingestion even at high doses is generally not considered a risk.⁴¹

Overall, none of the ingredients of this supplement are present at levels that pose any apparent risk to healthy, nonpregnant adults. On the other hand, supplementation with dietary minerals and vitamins that significantly exceed the recommended daily intakes carries with it an unknown risk, and therefore the long-term safety of the nutritional supplement used in this study cannot be defini-

tively proven. Patients undergoing clinical trials of this supplement should be monitored for adverse reactions and general health, just as they are when undergoing a trial of a new psychotropic medication.

Mechanisms of Action

An important question to address is *why* a broad-based nutrient supplement might improve mental health. There are at least 2 issues to consider:

1. Nutrient deficiencies influencing brain function. That there is a genetic predisposition to bipolar disorder is fairly well accepted.⁴² The question is, what do those genes do? It is possible that bipolar disorder is an inborn error of metabolism, analogous to others such as phenylketonuria in which metabolic "errors" lead to altered brain function, but whose symptoms become clinically evident long after birth. If this is the case, the fact that a nutritional supplement may partially correct that metabolic error suggests that the predisposing genes are coding for proteins involved in metabolic pathways dependent on some of those nutrients. Many minerals (e.g., zinc) are important in dozens of biochemical pathways vital to brain function, so this observation provokes many questions about the specific mechanisms by which predisposing genes might affect mental health.
2. Nutrient deficiencies in the food supply. In view of the wide range of biological variability in humans, it is certainly feasible that different individuals have variable vulnerabilities to diverse nutritional deficiencies. An amount of dietary mineral intake that is sufficient for most of the population may be borderline-adequate, deficient, or even toxic for a minority.

Although the claim that the nutrient content of our food supply is decreasing is often made but not substantiated by health food publications, there is actually a small amount of valid scientific data supporting this contention. Mayer⁴³ compared the mineral content of 20 fruits and 20 vegetables from 1936 to the 1980s using a laborious methodology to ensure that comparable laboratory methods were employed. Mayer's results were sobering: over that 50-year period, there were statistically significant decreases of calcium, magnesium, copper, and sodium in vegetables and of magnesium, iron, copper, and potassium in fruits. Zinc was not studied. The magnitude of some changes was large: for instance, the copper level in vegetables in the 1980s was less than 20% of the 1936 levels. As Mayer pointed out, the changes should not be surprising, as agriculture relies on fertilizers containing only nitrogen, phosphorous, and potassium, and there is little effort to remineralize the soil over the decades.

CONCLUSION

Regardless of the mechanisms involved, the findings reported here suggest that the amelioration of mood instability with broad spectrum nutritional interventions warrants further exploration.

Drug names: alprazolam (Xanax and others), amitriptyline (Elavil and others), benzotropine (Cogentin and others), bupropion (Wellbutrin and others), carbamazepine (Tegretol, Carbatrol), chlorpromazine (Thorazine and others), citalopram (Celexa), clomipramine (Anafranil and others), clonazepam (Klonopin and others), clonidine (Catapres and others), clozapine (Clozaril and others), desipramine (Norpramin and others), dextroamphetamine (Dexedrine and others), diazepam (Valium and others), doxepin (Sinequan and others), fluoxetine (Prozac and others), fluvoxamine (Luvox), gabapentin (Neurontin), haloperidol (Haldol and others), lamotrigine (Lamictal), levothyroxine (Synthroid and others), lorazepam (Ativan and others), methylphenidate (Ritalin and others), nefazodone (Serzone), nortriptyline (Pamelor and others), olanzapine (Zyprexa), paroxetine (Paxil), pemoline (Cylert), propranolol (Inderal and others), quetiapine (Seroquel), risperidone (Risperdal), sertraline (Zoloft), topiramate (Topamax), trazodone (Desyrel and others), trifluoperazine (Stelazine and others), trimipramine (Surmontil), valproic acid (Depakene and others), venlafaxine (Effexor), zolpidem (Ambien).

REFERENCES

- Alpert JE, Mischoulon D, Nierenberg AA, et al. Nutrition and depression: focus on folate. *Nutrition* 2000;16:544–546
- Fava M, Borus JS, Alpert JE, et al. Folate, vitamin B12, and homocysteine in major depressive disorder. *Am J Psychiatry* 1997;154:426–428
- Howell GA, Welch MG, Frederickson CJ. Stimulation-induced uptake and release of zinc in hippocampal slices. *Nature* 1984;308:736–738
- Dyck RH, Cynader MS. Histochemical localization of synaptic zinc in the developing kitten visual cortex. *J Comp Neurol* 1993;329:53–67
- Sandstead HH, Frederickson CJ, Penland JG. History of zinc as related to brain function. *J Nutr* 2000;130:496S–502S
- Vlachova V, Zemkova H, Vyklicky L Jr. Copper modulation of NMDA responses in mouse and rat cultured hippocampal neurons. *Eur J Neurosci* 1996;8:2257–2264
- Olivares M, Uauy R. Copper as an essential nutrient. *Am J Clin Nutr* 1996;63:791S–796S
- Hollmann M, Heinemann S. Cloned glutamate receptors. *Annu Rev Neurosci* 1994;17:31–108
- Martin SJ, Grimwood PD, Morris RGM. Synaptic plasticity and memory: an evaluation of the hypothesis. *Annu Rev Neurosci* 2000;23:649–711
- Caddell JL, Graziani LJ, Wiswell TE, et al. The possible role of magnesium in protection of premature infants from neurological syndromes and visual impairments and a review of survival of magnesium-exposed premature infants. *Magnes Res* 1999;12:201–216
- Maes M, D'Haese PC, Scharpe S, et al. Hypozincemia in depression. *J Affect Disord* 1994;31:135–140
- Maes M, Vandoolaeghe E, Neels H, et al. Lower serum zinc in major depression is a sensitive marker of treatment resistance and of the immune/inflammatory response in that illness. *Biol Psychiatry* 1997;42:349–358
- Walsh WJ, Isaacson HR, Rehman F, et al. Elevated blood copper/zinc ratios in assaultive young males. *Physiol Behav* 1997;62:327–329
- Toren P, Eldar S, Sela B-A, et al. Zinc deficiency in attention-deficit hyperactivity disorder. *Biol Psychiatry* 1996;40:1308–1310
- Arnold LE, Votolato NA, Kleykamp D, et al. Does hair zinc predict amphetamine improvement of ADD/hyperactivity? *Int J Neurosci* 1990;50:103–107
- Arnold LE, Pinkham SM, Votolato N. Does zinc moderate essential fatty acid and amphetamine treatment of attention-deficit/hyperactivity disorder? *J Child Adolesc Psychopharmacol* 2000;10:111–117
- Linder J, Brimar K, Granberg PO, et al. Characteristic changes in psychiatric symptoms, cortisol and melatonin but not prolactin in primary hyperparathyroidism. *Acta Psychiatr Scand* 1988;78:32–40
- Okamoto T, Gerstein HC, Obara T. Psychiatric symptoms, bone density and non-specific symptoms in patients with mild hypercalcemia due to primary hyperparathyroidism: a systematic overview of the literature. *Endocrinology* 1997;44:367–374
- Helmeste DM, Tang SW. The role of calcium in the etiology of the affective disorders. *Jap J Pharmacol* 1998;77:107–116
- Michelson D, Stratakis C, Hill L, et al. Bone mineral density in women with depression. *N Engl J Med* 1996;335:1176–1181
- Coelho R, Silva C, Maia A, et al. Bone mineral density and depression: a community study of women. *J Psychosom Res* 1999;46:29–35
- Benton D, Donohoe RT. The effects of nutrients on mood. *Public Health Nutr* 1999;2:403–409
- Hurtado EK, Claussen AH, Scott KG. Early childhood anemia and mild or moderate mental retardation. *Am J Clin Nutr* 1999;69:115–119
- Nelson C, Erikson K, Pinerio DJ, et al. In vivo dopamine metabolism is altered in iron-deficient anemic rats. *J Nutr* 1997;127:2282–2288
- Weiser M, Levkowitz Y, Neuman M, et al. Decrease of serum iron in acutely psychotic schizophrenic patients. *Int J Neurosci* 1994;78:49–52
- Levine J, Rapoport A, Mashiah M, et al. Serum and cerebrospinal levels of calcium and magnesium in acute versus remitted schizophrenic patients. *Neuropsychobiology* 1996;33:169–172
- Schaefer AL, Jones SDM, Stanley RW. The use of electrolyte solutions for reducing transport stress. *J Anim Sci* 1997;75:258–265
- First MB, Spitzer RL, Gibbon M, et al. Structured Clinical Interview for DSM-IV Axis I Disorders (SCID-I). Washington, DC: American Psychiatric Press; 1997
- Hamilton M. A rating scale for depression. *J Neurol Neurosurg Psychiatry* 1960;23:56–62
- Young RC, Biggs JT, Zieler VE, et al. A rating scale for mania: reliability, validity and sensitivity. *Br J Psychiatry* 1978;133:429–435
- Overall JE, Gorham DR. The Brief Psychiatric Rating Scale. *Psychol Rep* 1962;10:799–812
- Calabrese JR, Suppes T, Bowden CL, et al. A double-blind, placebo-controlled, prophylaxis study of lamotrigine in rapid-cycling bipolar disorder. *J Clin Psychiatry* 2000;61:841–850
- Kowatch RA, Suppes T, Carmody TJ, et al. Effect size of lithium, divalproex sodium, and carbamazepine in children and adolescents with bipolar disorder. *J Am Acad Child Adolesc Psychiatry* 2000;39:713–720
- Tohen M, Jacobs TG, Grundy SL, et al. Efficacy of olanzapine in acute bipolar mania: a double-blind, placebo-controlled study. *Arch Gen Psychiatry* 2000;57:841–849
- Hoffer A. Vitamin B-3 and Schizophrenia. Kingston, Ontario: Quarry Press; 1999:160
- Stoll AL, Severus E, Freeman MP, et al. Omega 3 fatty acids in bipolar disorder: a preliminary double-blind, placebo-controlled trial. *Arch Gen Psychiatry* 1999;56:407–412
- Meyers DG, Maloley PA, Weeks D. Safety of antioxidant vitamins. *Arch Intern Med* 1996;156:925–935
- Miller DR, Hayes KC. Vitamin excess and toxicity. In: Hathcock JN, ed. *Nutritional Toxicology*. New York, NY: Academic Press; 1982:81–116
- Omaye ST. Safety of megavitamin therapy. *Adv Exp Med Biol* 1984;177:169–203
- Yang G, Ge K, Chen J, et al. Selenium-related endemic diseases and the daily selenium requirement of humans. *World Rev Nutr Diet* 1988;22:98–152
- Hunt CD, Stoecker BJ. Deliberations and evaluations of the approaches, endpoints and paradigms for boron, chromium and fluoride dietary recommendations. *J Nutr* 1996;126(9 suppl):2441S–2451S
- Berrettini W. Diagnostic and genetic issues of depression and bipolar illness. *Pharmacotherapy* 1995;15:69S–75S
- Mayer AB. Historical changes in the mineral content of fruits and vegetables. *Br Food J* 1997;99:207–211

Appendix 1 appears on page 944.

Appendix 1. Supplement Facts: E.M.Power+ (serving size = 8 capsules, taken 4 times per day for study protocol)^a

| Ingredient | Amount Per Serving | % Daily Value |
|--|-----------------------|------------------|
| Vitamin A (as retinyl palmitate) | 3333 IU | 67 |
| Vitamin C (as ascorbic acid) | 250 mg | 417 |
| Vitamin D (as cholecalciferol) | 400 IU | 100 |
| Vitamin E (as <i>d</i> -alpha tocopheryl succinate) | 100 IU | 333 |
| Vitamin B ₁ (as thiamine mononitrate) | 5 mg | 333 |
| Vitamin B ₂ (as riboflavin) | 5.5 mg | 324 |
| Vitamin B ₃ (as niacinamide) | 25 mg | 125 |
| Vitamin B ₆ (as pyridoxine hydrochloride) | 7 mg | 350 |
| Vitamin B ₉ (as folic acid) | 400 µg | 100 |
| Vitamin B ₁₂ (as cyanocobalamin) | 250 µg | 4167 |
| Biotin | 25 µg | 8 |
| Pantothenic acid (as <i>d</i> -calcium pantothenate) | 6 mg | 60 |
| Calcium (as calcium complex,* calcium amino acid chelate) | 550 mg | 55 |
| Iron (as iron amino chelate, iron complex*) | 6 mg | 33 |
| Phosphorous (phosphorous complex) | 350 mg | 35 |
| Iodine (from kelp) | 75 µg | 50 |
| Magnesium (as magnesium amino acid chelate, magnesium complex) | 250 mg | 63 |
| Zinc (as zinc amino acid chelate, zinc complex*) | 20 mg | 133 |
| Selenium (as selenium amino acid chelate, selenium complex*) | 100 µg | 143 |
| Copper (as copper amino acid chelate, copper complex*) | 3 mg | 150 |
| Manganese (as manganese amino acid chelate, manganese complex) | 4 mg | 200 |
| Chromium (as chromium amino acid chelate, chromium complex*) | 250 µg | 208 |
| Molybdenum (as molybdenum amino acid chelate, molybdenum complex) | 66 µg | 88 |
| Potassium (as potassium complex*) | 100 mg | 3 |
| And a proprietary blend of the following**: | | |
| <i>dl</i> -Phenylalanine, glutamine (as <i>l</i> -glutamine), citrus bioflavonoids (from peel), grape seed (<i>Vitis vinifera</i>), choline (as choline bitartrate), inositol, <i>Ginkgo biloba</i> (from leaf), methionine (as <i>l</i> -methionine), germanium (as <i>Germanium sesquioxide</i>), boron (as boron amino acid chelate), vanadium (as vanadium amino acid chelate, vanadium complex*), nickel (as nickel amino acid chelate, nickel complex*) | | |
| Other ingredients: gelatin, magnesium stearate, microcrystalline cellulose, silicon dioxide | | |

^aManufactured for The Synergy Group of Canada, by Evinco International, Farmington, Utah.

*Saccharide complex.

**Daily value not established.